

Heuristic analysis of the Pure Random Walk search procedure for random SAT problems

(on the importance of large deviations for the...)

Guilhem Semerjian

Ecole Normale Supérieure, Paris

Rémi Monasson

Ecole Normale Supérieure, Paris

Univ. Louis Pasteur, Strasbourg

Collaborator :

Simona Cocco

Univ. Louis Pasteur, Strasbourg

Pure Random Walk procedure for 3-SAT

1. Pick up an initial (random) configuration of variables;
2. If all clauses are satisfied, **STOP**.
3. Else, choose randomly one unsat clause and flip one of its attached variables;
Goto 2.

Example:

set of	(NOT x_1 or x_2 or x_3)	→	F
clauses	(x_1 or x_2 or NOT x_4)	→	T
	(NOT x_1 or x_2 or x_4)	→	T

1. random configuration $x_1=T, x_2=F, x_3=F, x_4=T$
2. pick up first clause and flip one variable it contains
e.g. $x_2=F$ $\ddot{\circ}$ $x_2=T$ then all clauses are satisfied
e.g. $x_1=T$ $\ddot{\circ}$ $x_1=F$ then clause 1 is satisfied, but clause 2 is not any longer.

Number of unsat clauses (=energy) is not guaranteed to decrease!

Rigorous results for 2-Sat and 3-Sat

- Papadimitriou (1992)

2-SAT: if there is a solution, it is found w.h.p. in a time growing as $O(N^2)$.

- Schönning (2000):

3-SAT: the probability that a satisfiable instance is not solved after T runs (of $3N$ steps each) is bounded from above by $\exp(-T\mu(3/4)^N)$
recently: 1.329... instead of 1.333...


- Alekhnovich and Ben-Sasson (2002):

random 3-SAT: a solution is almost surely reached in polynomial time if ratio a of clauses/variable is smaller than 1.63


Phenomenology

Single runs on instances with $N=500$ variables, $M= a N$ clauses

$a = 2$



$a = 3$



(Approximate) Analysis

- **Polynomial** regime: what is the average solving time?

$$T_{\text{res}}(N, a) = N t_{\text{res}}(a)$$

- Characterization of the **polynomial/exponential** transition.
(average behaviour: plateau properties)

- **Exponential** regime: what is the average solving time?

$$\begin{aligned} T_{\text{res}}(N, a) &= \exp(N z(a)) \\ &= 1 / \text{Proba}(j_0 = 0) = \exp(M p(0)) \end{aligned}$$

(large deviations: escape from plateau)

Average running time in the polynomial regime

- if α is small enough, the graph of interactions is essentially made of small clusters C
- $t_{\text{res}} = \sum_C p(C) t_{\text{res}}(C)$
- results for K-SAT (*exact provided t_{res} can be expanded*):

$$t_{\text{res}}(\alpha, K) = \frac{1}{2^K} + \frac{K(K+1)}{K-1} \frac{1}{2^{2K+1}} \alpha + \frac{4K^6 + K^5 + 6K^3 - 10K^2 + 2K}{3(K-1)(2K-1)(K^2-2)} \frac{1}{2^{3K+1}} \alpha^2 + O(\alpha^3)$$

But cannot capture the plateau regime ...

Average behaviour (I)

Parameters:

N = number of variables

M = number of clauses = $a N$

K = number of variables in each clause (K-SAT)

- Define $M_j(T)$ the number of **clauses satisfied by j literals** after T clause flips with $j = 0, 1, \dots, K$.
($M_0(T)$ is the number of unsatisfied clauses)

- Normalization condition:
$$\sum_{j=0}^K M_j(T) = M$$


- Define fractions of **j -clauses**:
$$\phi_j(t) = M_j(T)/M$$

Question: evolution equations for the fractions when $T \rightarrow T+1$?

Average behaviour (II)

Initial conditions for the fractions of j -clauses: $\phi_j(0) = \frac{1}{2^K} \binom{K}{j}$

After a new clause flip, i.e. from $T \rightarrow T+1$:



$$M_j(T+1) - M_j(T) = \frac{K}{N} \left(\frac{(K-j+1)}{K} M_{j-1}(T) + \frac{(j+1)}{K} M_{j+1}(T) \right) - d_j + d_{j-1}$$

Average behaviour (III)

DT = 1 leads to $DM_j = \alpha(1)$. Thus, fractions F_j vary by $\alpha(1)$ on time scale DT = $\alpha(M)$. Define reduced time: $t = T/M$.

Evolution equation:
$$\frac{d\vec{\phi}}{dt} = \vec{v} + \alpha K W_r \cdot \vec{\phi} \quad \text{with}$$

$$\vec{v} = \begin{pmatrix} -1 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \quad W_r = \begin{pmatrix} -1 & \frac{1}{K} & 0 & 0 & 0 & \dots & 0 \\ 1 & -1 & \frac{2}{K} & 0 & 0 & \dots & 0 \\ 0 & \frac{K-1}{K} & -1 & \frac{3}{K} & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & \dots & -1 \end{pmatrix}$$

NB: normalization of fractions comes from nullity of matrix elements summed over columns.

Average behaviour (IV)

Existence of a critical value of the ratio clauses/variable


$$\alpha_d(K) = \frac{2^K - 1}{K}$$

- If $a < a_d$, then the procedure stops after a finite time with no unsat clause left.

$$t_{\text{res}}(a, K) = \frac{1}{a} \operatorname{Tanh}^{-1} \left[\left(1 - a/a_d(K) \right)^{-1/K} - 1 \right]$$

- If $a > a_d$, then the procedure never stops, and a finite fraction of clauses remain unsatisfied.

$$\phi_0^* = \frac{1}{2^K} - \frac{1}{\alpha K} \left(1 - \frac{1}{2^K} \right) = \frac{1}{2^K} \left(1 - \frac{\alpha_d(K)}{\alpha} \right)$$



Large deviations and escape from plateau (I)

Evolution equation for the vector of clause types populations:

$$\text{Prob}[\mathbf{M}', T + 1] = \sum_{\mathbf{M}} [A_{\mathbf{M}'\mathbf{M}}] \text{Prob}[\mathbf{M}, T] \quad ,$$

where

$$[A_{\mathbf{M}'\mathbf{M}}] = \sum_{\mathbf{Z}} \frac{N Z_0}{K M_0} \delta(\mathbf{M}' - \mathbf{M} - \Delta \cdot \mathbf{Z}) P(\mathbf{Z}|\mathbf{M})$$

Z_i = number of clauses of type i in which the flipped variable appears,

Z_i^s = number of clauses of type i in which the flipped variable was one of the i satisfying literal.

$$P(\mathbf{Z}|\mathbf{M}) = \prod_{i=0}^K \binom{M_i}{Z_i} \left(\frac{K}{N}\right)^{Z_i} \left(1 - \frac{K}{N}\right)^{M_i - Z_i} \prod_{i=1}^{K-1} \binom{Z_i}{Z_i^s} \left(\frac{i}{K}\right)^{Z_i^s} \left(1 - \frac{i}{K}\right)^{Z_i - Z_i^s}$$

Large deviations and escape from plateau (II)

Scaling hypothesis for clause type populations:

$$\text{Prob}[\mathbf{M}, \mathbf{T}] \propto \exp(-M \rho(\mathbf{j}, t)) \quad \text{where} \quad \begin{aligned} j_j &= \mathbf{M}_j / M \\ t &= \mathbf{T} / M \end{aligned}$$

Resulting partial differential equation for ρ :

$$\rho_t = -\rho_0 + \rho_1 + a \sum_{j=0}^K j_j \left((K-j) e^{\rho_{j+1} - \rho_j} + j e^{\rho_{j-1} - \rho_j} - K \right)$$

with initial condition:


$$\rho(\mathbf{j}, t=0) = \sum_{j=0}^K j_j \ln(j_j / j_j^0) \quad ; \quad j_j^0 = \frac{1}{2^K} \binom{K}{j}$$

Large deviations and escape from plateau (III)

upper
bound = $\ln 4/3$




$$z = p(0)/a$$




Intuitive summary: the rolling ball picture

On the plateau: $P_N(j_0) \approx \exp(-M p(j_0))$



Exponential regime



Polynomial regime

Conclusions (I)

1. « Approximate » analysis of Pure Random Walk for random SAT problems (also XOR-SAT)

- Annealed approximation may be justified for infinite a, K at fixed $a^* = aK/2^K$

$$a_d^* = 1 \quad (\text{and } a_c^* = \frac{1}{2})$$

$$t_{res}^*(\alpha^*) = -\frac{1}{\alpha^*} \ln(1 - \alpha^*) = 1 + \frac{1}{2} \alpha^* + \frac{1}{3} (\alpha^*)^2 + O((\alpha^*)^3)$$

*Approximate
result*

*Exact
expansion*

- Finite K corrections ? Much tougher, under way ...
- What can be said from a rigorous point of view?

Conclusions (II)

2. Exponentially rare events (large deviations) matter for analyzing algorithms

- notion of metastability
- case of DPLL and random restarts

3. Technical set-up:

Average behaviour

Ordinary Differential Equations

Large deviations

Partial Differential Equations