

ICFP M2 - STATISTICAL PHYSICS 2
 Solution of the homework n° 5
 Langevin and Fokker-Planck equations

Guilhem Semerjian

March 2019

1. The Fokker-Planck equation is indeed written here as a continuity equation that traduces the local conservation of the probability (or of the number of particles if we don't normalize P to 1). The probability $P(x, t)dx$ of finding the particle in the infinitesimal interval $[x, x + dx]$ evolves during the infinitesimal time dt because of the incoming flux $J(x, t)dt$ and of the outgoing flux $-J(x + dx, t)dt$. One can thus interpret $J(x, t)$ as the current of particles flowing through the position x at time t , counted in the increasing x direction. The first contribution to J arises from the deterministic force $-V'(x)$, and counts the number of particles that crosses x with this velocity. The second term is a diffusion effect, the random force η tends to equalize the density of presence of the particle.
2. Δx is a sum of Gaussian random variables, it is thus Gaussian. One can characterize it by its two first moments,

$$\mathbb{E}[\Delta x] = \int_t^{t+\Delta t} dt' \mathbb{E}[\eta(t')] = 0, \quad (1)$$

$$\mathbb{E}[(\Delta x)^2] = \int_t^{t+\Delta t} dt'_1 \int_t^{t+\Delta t} dt'_2 \mathbb{E}[\eta(t'_1)\eta(t'_2)] = 2T\Delta t. \quad (2)$$

3. (a) When $T = 0$ the random term disappears from the Langevin equation, the deterministic trajectory $x_*(t)$ of the particle is thus the solution of the ordinary differential equation $x'_*(t) = -V'(x_*(t))$ with the initial condition $x_*(t = 0) = x_0$. The solution of the Fokker-Planck equation is then $P(x, t) = \delta(x - x_*(t))$.
- (b) When $V(x)$ is independent of x there is no deterministic force and the Fokker-Planck equation reduces to the diffusion equation $\frac{\partial P}{\partial t} = T \frac{\partial^2 P}{\partial x^2}$. The solution of this equation with the initial condition peaked in x_0 is the Gaussian distribution

$$P(x, t) = \frac{1}{\sqrt{4\pi Tt}} e^{-\frac{(x-x_0)^2}{4Tt}}. \quad (3)$$

The solution of the Langevin equation is

$$x(t) = x_0 + \int_0^t dt' \eta(t'), \quad (4)$$

which is indeed a Gaussian random variable centered in x_0 with variance $2Tt$.

4. One has $T \frac{dP_{\text{GB}}}{dx} = -V'(x)P_{\text{GB}}(x)$, hence the current J vanishes for this choice of P , which makes $P_{\text{GB}}(x)$ a stationary solution of the Fokker-Planck distribution. If the potential $V(x)$ is confining, in such a way that the Gibbs-Boltzmann distribution is normalizable, one has $P(x, t) \rightarrow P_{\text{GB}}(x)$ at large times, for all initial conditions.

5. In this case the Langevin equation $\frac{dx}{dt} = -x(t) + \eta(t)$ can be integrated with, for instance, the variation of constant method, to give

$$x(t) = x_0 e^{-t} + \int_0^t dt' e^{-(t-t')} \eta(t') . \quad (5)$$

The mean of this Gaussian random variable is $x_0 e^{-t}$, that decays to zero at large time : the particle forgets the initial condition and relaxes on average to the bottom of the potential well. The variance of $x(t)$ reads

$$\int_0^t dt'_1 \int_0^t dt'_2 e^{-(2t-t'_1-t'_2)} \mathbb{E}[\eta(t'_1)\eta(t'_2)] = 2T \int_0^t dt' e^{-2(t-t')} = T(1 - e^{-2t}) , \quad (6)$$

it grows with time towards the equilibrium one.