

Contents

1 General Introduction, Basic Concepts and Tools: Quantum Statistics and Interaction	1
1.1 The Ideal Bose Gas: Reminders and Appetizers	3
1.1.1 In the Grand Canonical Ensemble	4
1.1.1.1 Condensation by saturation of excited modes . . .	5
1.1.1.2 One-body signatures: density profile, g_1 function .	8
1.1.1.3 Two-body signatures: g_2 function, giant fluctuations of n_0 , symmetric partition noise	12
1.1.2 In the canonical then microcanonical ensemble	14
1.1.2.1 Experimental motivation	14
1.1.2.2 Elimination of the condensate mode and the never-empty condensate approximation	15
1.1.2.3 Canonical fluctuations of $\hat{N}_0$: first moments and probability distribution	16
1.1.2.4 Microcanonical fluctuations of $\hat{N}_0$	20
1.2 What model for the interaction	22
1.2.1 The problem of metastability, the solution by universality and the discretization of space	22
1.2.2 The T matrix as a guiding thread	27
1.2.3 The notion of s -wave scattering length	29
1.2.3.1 In three dimensions	29
1.2.3.2 In reduced dimensionality	31
1.3 Scattering amplitude, T matrix and link with cold atoms for the Wigner-Bethe-Peierls model of a zero-range interaction	32
1.3.1 In three dimensions	32
1.3.2 In reduced dimensionality	37
1.3.2.1 Case $d = 2$	37
1.3.2.2 Case $d = 1$	41
1.4 The Simplest Contact Interaction: The Lattice Model	43
1.4.1 In Discretized Real Space	43
1.4.2 In Momentum Space	45
1.4.3 T matrix, bare coupling constant and Hamiltonian	46

2 The pure condensate regime: Gross-Pitaevskii equation	49
2.1 Stationary case	52
2.1.1 In three dimensions	52
2.1.1.1 A variational formulation	52
2.1.1.2 Case $a > 0$ and healing length ξ	53
2.1.1.3 Case $a < 0$ and collapse instability	54
2.1.1.4 Trapped case and Thomas-Fermi limit	54
2.1.2 In two dimensions	57
2.1.2.1 A questionable scale invariance	57
2.1.2.2 Thomas-Fermi limit	59
2.1.3 In one dimension: the bright soliton	59
2.1.3.1 Critical analysis of the coupling constant	61
2.1.3.2 Applications: equation of state, Thomas-Fermi limit, bright soliton and breaking of translational invariance	62
2.1.3.3 What we learn from the Bethe ansatz: high or low density, condensate or not, quantum soliton, Schrödinger's cat, liquid-gas transition	64
2.1.4 Complement: local energy minimization condition	69
2.2 Applications: stationary condensates with phase defects	73
2.2.1 An equation enriched by a rotation term	73
2.2.2 In one dimension: gray soliton and second Lieb branch	76
2.2.2.1 A saving formulation	76
2.2.2.2 Integrability of the nonlinear Schrödinger equation	78
2.2.2.3 In the thermodynamic limit	79
2.2.2.4 Obtaining the second Lieb branch by energy	80
2.2.2.5 Obtaining the second Lieb branch by phase shift	81
2.2.2.6 And in the attractive case?	83
2.2.3 In two dimensions: rotating trapped condensates with quantum vortices	84
2.2.3.1 A Thomas-Fermi trial wavefunction with vortices	85
2.2.3.2 Mean energy of n vortices	87
2.2.3.3 Physical discussion at fixed Ω	89
2.2.3.4 Physical discussion at fixed L_z for $n = 1$	92
2.2.3.5 Neither condensates in the strict sense nor superfluids	93
2.2.4 In three dimensions: the curved vortex line	95
2.2.4.1 A simple reasoning by cutting into slices	95
2.2.4.2 Numerically tested predictions	97
2.2.4.3 At fixed angular momentum L_z	98
2.3 Time-dependent case	99
2.3.1 Form of the equation and its dimensional reductions	101

2.3.2	Hydrodynamic equations as an equivalent of the Thomas-Fermi approximation in the time-dependent case, and how to solve them	102
2.3.2.1	Obtaining by switching to phase-modulus representation	102
2.3.2.2	Solution of the hydrodynamic equations	104
2.3.2.3	For a Lagrangian specification of the flow field	106
2.3.2.4	Linearized hydrodynamic equations	107
2.3.2.5	Eigenmodes in the absence of rotation	109
2.3.3	Some exact solutions of the Gross-Pitaevskii equation	115
2.3.3.1	1D case: setting the bright soliton in motion	115
2.3.3.2	2D case: reducing to a trap of constant stiffness by changing gauge and scale	117
2.3.3.3	General case: modifying the overall motion in a trap	121
2.3.4	Application: elucidating the mechanism of vortex lattice formation in the ENS experiment	122
2.3.4.1	The ENS experimental procedure and the failure of thermodynamic scenarios	122
2.3.4.2	A new two-step mechanism, resonance and dynamic instability	125
2.3.4.3	A verification procedure: slow rotation	129
2.3.4.4	The Landau thermodynamic scenario, continued and concluded	134
2.3.4.5	Moral	136
2.3.4.6	Complement: General stationary solutions of the hydrodynamic equations in a harmonic trap rotating around one of its eigenaxes	137
2.3.5	Study of dynamic stability	140
2.3.5.1	A very simple calculation	141
2.3.5.2	A more rigorous derivation of $\mathcal{L}(t)$	143
2.3.5.3	Time-independent case: normal and anomalous eigenmodes, Bogoliubov transformation, dynamic and thermodynamic stability	144
2.3.5.4	Time-dependent case	153
2.4	Limitations to the validity of the Gross-Pitaevskii equation for the time evolution	154
2.4.1	Evolution of a bright soliton « at rest »	155
2.4.1.1	By Heisenberg equations for the quantum field	156
2.4.1.2	By linear stability analysis for the classical field	157
2.4.2	A phase scrambling mechanism omitted by the Gross-Pitaevskii equation	159
2.4.2.1	Two-Mode Model	159

2.4.2.2	Classical Field versus Quantum Field	160
2.4.2.3	Improvement of the Classical Field by Adding Wigner Noise to the initial State	163
2.4.3	Generalities of this scrambling mechanism: breathing mode in an isotropic harmonic trap	164
2.4.3.1	Excitation by change in trap stiffness	164
2.4.3.2	Stability of the breathing mode: effect of scale fac- tor fluctuations	166
2.4.3.3	Stability of other modes	169
2.4.4	Absence of the spontaneous emission mechanism in the Gross-Pitaevskii equation	170
2.4.4.1	An analogy with quantum radiation	170
2.4.4.2	Fast quantum depletion in a two-mode model	171
2.4.4.3	In classical field improved by Wigner noise	175
2.4.5	Another, multimode example dominated by spontaneous pair emission: twin beams	176
2.4.5.1	1D model with time-modulated coupling constant	176
2.4.5.2	Linear stability analysis for classical fields	178
2.4.5.3	Statistical properties of twin beams	181
2.4.6	Moral of the discussion on the validity of Gross-Pitaevskii	182
3	Bogoliubov's theory: first corrections to the pure condensate in dimen- sion three and the condensate phase operator	183
3.1	General idea of the Bogoliubov method	184
3.1.1	The non-condensed field $\hat{\psi}_\perp$ as a perturbation	184
3.1.2	Implementation: Hamiltonian expansion, elimination of the condensate mode, phase operator $\hat{\theta}$ and redefined non- condensed field $\hat{\Lambda}$, correction to Gross-Pitaevskii	185
3.1.3	A historic breakthrough	188
3.2	Spatially homogeneous stationary case	189
3.2.1	The lattice model case	190
3.2.1.1	Implementation of the Bogoliubov method and first physical interpretations	190
3.2.1.2	A hidden expansion	193
3.2.1.3	Final form of Bogoliubov Hamiltonian; excitation spectrum, ground state energy	195
3.2.2	For a true interaction potential $V(\mathbf{r})$	199
3.2.3	Simple applications: statistics of n_0 , anomalous noncon- densed density, equation of state, momentum distribution and correlations, g_1 and g_2 functions, time coherence of the noncondensed field	202
3.2.3.1	Statistics of n_0 at equilibrium	202

3.2.3.2	The anomalous noncondensed density ρ_{an}	210
3.2.3.3	The equation of state of the weakly interacting Bose gas in Bogoliubov approximation	211
3.2.3.4	First-order coherence function g_1 and wavevector distribution $n_{\mathbf{k}}^{\text{cin}}$ of the gas; correlation function in momentum space	221
3.2.3.5	Pair distribution function g_2	225
3.2.3.6	Spatio-temporal coherence function g_1 in the Bogoliubov approximation	229
3.2.4	The special case of superfluidity	237
3.2.4.1	Superfluidity is not Condensation	237
3.2.4.2	Landau Critical Velocity and Beyond	238
3.2.4.3	Metastable Currents and Their Bogoliubov Analysis	242
3.2.4.4	Thermodynamic Definition of the Normal Fraction	249
3.2.4.5	Energy-based Variant and Leggett Bound	252
3.2.5	Complement I: Presentation and implementation on the normal and anomalous noncondensed densities of a general high- and low-temperature expansion method, and challenging the Hartree-Fock theory	256
3.2.5.1	The noncondensed density	256
3.2.5.2	At leading order in temperature	258
3.2.5.3	How to go beyond leading order	259
3.2.5.4	The anomalous noncondensed density	262
3.2.5.5	What is the point of high-temperature expansion?	264
3.2.5.6	Application: challenging Hartree-Fock	264
3.2.6	Complement II: Quantum adiabaticity and thermodynamic adiabaticity	267
3.3	Stationary case in a trap	271
3.3.1	Motivation and specificities	271
3.3.1.1	Which spatial mode of the condensate?	271
3.3.1.2	Which Bogoliubov modes? Semi-classical limit . .	272
3.3.2	A simplified case to understand why order-3 terms in $f_{\text{nc}}^{1/2}$ in the Hamiltonian can affect mean values at order 2	274
3.3.2.1	Bogoliubov calculation for a single degree of freedom	274
3.3.2.2	Moral	277
3.3.3	Cubic Approximation of the Hamiltonian	277
3.3.3.1	First Step of Cubization	277
3.3.3.2	Second Step of Cubization	278
3.3.3.3	Final Result and Its Interpretation	279
3.3.4	Relationship between $\hat{\Lambda}^{(2)}$, $\langle \hat{\Lambda}^{(1)} \rangle$ and $\phi_{\perp}^{(2)}$	282
3.3.5	Explicit expansion of the theory to order 3 in $f_{\text{nc}}^{1/2}$	283

3.3.5.1	Overview of the rest of the expansion in $f_{\text{nc}}^{1/2}$. . .	283
3.3.5.2	At order 0 in $f_{\text{nc}}^{1/2}$	283
3.3.5.3	At order 1 in $f_{\text{nc}}^{1/2}$	284
3.3.5.4	At order 2 in $f_{\text{nc}}^{1/2}$: the discrete Bogoliubov Hamiltonian and its reduced form	284
3.3.5.5	At order 3 in $f_{\text{nc}}^{1/2}$: condensate wave function beyond Gross-Pitaevskii	290
3.3.5.6	Calculation of $\langle \hat{\Lambda}^{(1)}(\mathbf{r}) \rangle_{\hat{H}^{(0-2)} + \hat{H}^{(3)}}^{(2)}$ and physical interpretation by depletion-interaction	292
3.3.6	Expansion of g_0 to order one in a/b and passage to the continuous limit (or negligible range interaction) $b/\xi \rightarrow 0$.	296
3.3.6.1	Context and motivation	296
3.3.6.2	Geometry considered for the continuous limit . .	297
3.3.6.3	Case of the ground state energy	298
3.3.6.4	Case of the condensate wave function	301
3.3.7	Summary: Direct formulation in continuous space of the time-independent Bogoliubov theory	303
3.3.7.1	Motivation and derivation	303
3.3.7.2	Case of $\phi^{(2)}$, first correction to Gross-Pitaevskii on the spatial mode of the condensate	305
3.3.7.3	Case of the Bogoliubov Hamiltonian and its ground energy level	308
3.3.7.4	Case of a more general observable	310
3.3.7.5	Continuous Space Summary: Main Steps and Equations	310
3.4	Bogoliubov Theory in the Time-Dependent Case	312
3.4.1	Physical Motivations and Overview	312
3.4.1.1	A Useful Question and a Fundamental Problem . .	312
3.4.1.2	A first step beyond Gross-Pitaevskii	313
3.4.1.3	A second step beyond Gross-Pitaevskii	314
3.4.1.4	Overview of the Method	315
3.4.2	Equation of Motion for the Field $\hat{\Lambda}$ to Order 2 in $f_{\text{nc}}^{1/2}$	315
3.4.3	Wavefunction of the condensate to order 0 in $f_{\text{nc}}^{1/2}$	321
3.4.4	Evolution of the non-condensed field and correction to $\phi^{(0)}$ to order 1 in $f_{\text{nc}}^{1/2}$	323
3.4.5	Contribution of order 2 in $f_{\text{nc}}^{1/2}$ to the wavefunction of the condensate	325
3.4.6	Expansion of g_0 to first order in a/b and continuous (or negligible range) limit $b/\xi \rightarrow 0$	327
3.4.7	Summary: direct formulation of the time-dependent Bogoliubov theory in continuous space	331
3.4.7.1	What small parameter?	332

3.4.7.2	Zeroth order: the Gross-Pitaevskii equation recovered	332
3.4.7.3	First order: modes of non-interacting quasiparticles	332
3.4.7.4	Second order: first correction to Gross-Pitaevskii	335
3.4.7.5	The importance of $\phi_{\perp}^{(2)}$ in observables	336
3.4.7.6	What if N fluctuates?	338
3.4.8	Connection with classical field physics and the truncated Wigner approximation	339
3.4.9	Different scenarios for exiting the validity regime of the time-dependent Gross-Pitaevskii equation	344
3.4.9.1	Failure by depletion divergence	345
3.4.9.2	Failure at the level of $\phi^{(2)}$	348
3.5	The condensate phase operator	349
3.5.1	Introduction of the phase operator $\hat{\theta}_{\phi}$	350
3.5.2	Evolution equation of $\hat{\theta}_{\phi}$ when $\partial_t \phi \equiv 0$	352
3.5.3	Temporal smoothing $\overline{d\hat{\theta}_{\phi}/dt}_t$ of the equation of evolution	354
3.5.3.1	Motivation and implementation	354
3.5.3.2	Smoothing quadratic terms in $d\hat{\theta}_{\phi}/dt$	356
3.5.3.3	Smoothing linear terms in $d\hat{\theta}_{\phi}/dt$: it is necessary to know the quadratic terms $\hat{S}^{(2)}$ of $d\hat{\Lambda}_{\phi^{(0)}}/dt$	357
3.5.3.4	Smoothing the source operator $\hat{S}^{(2)}$ and an operator $\hat{\phi}^{(2)}$	359
3.5.3.5	Smoothing linear terms in $d\hat{\theta}_{\phi}/dt$: continued and end	360
3.5.4	Recognizing a chemical potential operator in $d\hat{\theta}_{\phi}/dt$	362
3.5.4.1	Recognizing derivatives with respect to N	362
3.5.4.2	Manipulating Derivatives in Three Steps	364
3.5.4.3	Collecting terms, Final Result, and Interpretation	366
3.5.5	Beyond the Bogoliubov Approximation: The Spatially Homogeneous Case	367
3.5.5.1	Purpose, Idea, and Implementation of the Calculation	367
3.5.5.2	The result; its diagonal terms; its non-diagonal terms, phase diffusion and temporal smoothing	369
3.5.6	Case where the particle number fluctuates	371
3.5.6.1	Motivation	371
3.5.6.2	Extension of the calculation of $\overline{d\hat{\theta}_{\phi}/dt}_t$ from the previous sections	372
3.5.6.3	Further simplification for a large system	374
3.5.6.4	The result and its application to a statistical mixture of canonical ensembles	375

3.5.7	In the spatially homogeneous case, without temporal smoothing (with oscillating terms)	376
3.5.7.1	At fixed N	376
3.5.7.2	When N fluctuates	378
3.5.8	Complement: The paradoxes of the phase operator	379
4	Application I: Damping and energy shift of the excitation modes of a spatially homogeneous condensate	381
4.1	Obtaining by analysis of a Bragg excitation of low amplitude ϵ . . .	383
4.1.1	A thought experiment	383
4.1.2	Solution of the Gross-Pitaevskii equation at first order in ϵ .	385
4.1.3	Evolution of the Bogoliubov modes at first order in ϵ , and amplitudes $\mathcal{A}_{k_1, k_2}^{k_3}, \mathcal{A}_{k_1, k_2, k_3}$	387
4.1.4	Correction $\phi^{(2)}$ to the Gross-Pitaevskii wave function at first order in ϵ	390
4.1.4.1	Structure of the result and general considerations	390
4.1.4.2	Explicit calculation of $\phi^{(2)}$ at long times	393
4.1.5	Result: the first correction to Gross-Pitaevskii on the eigenfrequencies of the condensate	399
4.1.5.1	Three equivalent expressions	400
4.1.5.2	Overview of the literature on this topic	402
4.1.5.3	Conditions of applicability of result (4.84)	403
4.2	Obtaining and physical interpretation in terms of interaction between Bogoliubov quasiparticles	405
4.2.1	Bragg signal and correlation functions of the unperturbed system	405
4.2.2	Calculation by the quantum master equation and the quantum regression theorem	407
4.2.2.1	Interaction process between quasiparticles and recovered amplitudes $\mathcal{A}$	408
4.2.2.2	The quantum master equation in the Born-Markov approximation	410
4.2.2.3	The quantum regression theorem	412
4.2.3	Recovering the result of section 4.1.5	414
4.2.3.1	An apparent disagreement	414
4.2.3.2	Resolution by inclusion of $\hat{H}_4$	415
4.2.4	A validity condition of the result: that of the golden rule . .	417
4.3	Some explicit results on $\omega_{\mathbf{q}}^{(2)}$	419
4.3.1	Summary: all analytical results on $\omega_{\mathbf{q}}^{(2)}$ and numerical illustrations	419
4.3.1.1	Classification by running variable (tending to 0 or $+\infty$)	420

4.3.1.2	Numerical illustrations after reduction to a single integral	423
4.3.1.3	Complement: Calculation of integrals over λ in equations (4.160), (4.162) and (4.167)	429
4.3.2	Case $T = 0$: expansion of $\omega_{\mathbf{q}}^{(2)}$ at low q to order 4; the result is real but requires a multiscale expansion	429
4.3.2.1	Initial motivation: what speed of sound?	429
4.3.2.2	A too naive expansion under the integral sign	431
4.3.2.3	The correct method is multiscale	432
4.3.2.4	A smooth reconnection as a test	433
4.3.2.5	The final result to order 4	435
4.3.3	Case $T = 0$: expansion of $\omega_{\mathbf{q}}^{(2)}$ at low q to order 5; the result becomes complex and a logarithm of q appears	435
4.3.3.1	A real physical motivation	435
4.3.3.2	A model integral for practice, which shows the importance of the curvature of the dispersion relation	436
4.3.3.3	Back to the real problem	442
4.3.3.4	The result to order 5	447
4.3.4	Case $T \neq 0$: linearization of $\omega_{\mathbf{q}}^{(2)}$ at low q	447
4.3.4.1	Motivations and overview	447
4.3.4.2	Expansion in the area $k < Aq$	450
4.3.4.3	Expansion in the area $k > Aq$	452
4.3.4.4	A more pleasant final form	453
4.3.4.5	At low temperature	454
4.3.4.6	At high temperature	455
4.3.4.7	At any temperature	457
4.3.5	What is $\omega_{\mathbf{q}}^{(2)}$ for large q ?	458
4.3.5.1	Physical Motivation	458
4.3.5.2	What intuition suggests about the real part	459
4.3.5.3	What intuition suggests about the imaginary part	461
4.3.5.4	Expansion of $\omega_{\mathbf{q}}^{(2)}$ at large q at $T = 0$	463
4.3.5.5	At $T > 0$: large- q expansion of the thermal part of $\hbar\omega_{\mathbf{q}}^{(2)}$	468
4.3.6	Study of the thermal part $\omega_{\mathbf{q}}^{(2)\text{th}}$ of $\omega_{\mathbf{q}}^{(2)}$ at low and high temperatures	471
4.3.6.1	Low- T limit at fixed number wave number q	472
4.3.6.2	Low- T limit at fixed ratio $\hbar c_{\text{GP}} q / k_{\text{B}} T$	473
4.3.6.3	High- T limit at fixed wave number q	476
4.3.6.4	High- T limit at fixed ratio $\hbar^2 q^2 / m k_{\text{B}} T$	479
4.3.7	Complement I: Sectorization of Beliaev and Landau processes in wave vector space	482

4.3.8	Complement II: Boundary singularities of the integrand of $\int dk$ in $\omega_{\mathbf{q}}^{(2)}$	485
4.4	Moral of our calculation of $\omega_{\mathbf{q}}^{(2)}$	490
Main Notations		493
Index		499
Bibliography		511
5	Application II: Spin Squeezing and Its Limits; Schrödinger Cat States	537
5.1	Introduction	537
5.1.1	Principle of an atomic clock	537
5.1.2	Standard quantum noise	538
5.1.3	Squeezed States	540
5.1.4	Some Points for Further Study	540
5.1.5	Objectives of this Chapter	541
5.1.5.1	A Giant Spin Thanks to Condensates...	541
5.1.5.2	... but a multimode problem	542
5.2	Squeezing in the Kitagawa-Ueda model	543
5.2.1	Position of the Problem	543
5.2.2	Solution of the operator equations of motion	544
5.2.3	Bosonic representation and phase states	545
5.2.4	Mean spin, collapse and revival, Schrödinger's cat	547
5.2.5	Spin variances and covariances	548
5.2.6	Optimization of $\Delta S_{\perp, \min}$ over time	550
5.2.6.1	A first time scale	551
5.2.6.2	A second time scale time	552
5.2.7	Optimization of $\xi^2(t)$ over time	553
5.3	Effect of a stationary random dephasing environment	556
5.3.1	Motivation and presentation of the model	556
5.3.2	Larmor fluctuations and spin squeezing factor	557
5.3.3	What scaling law for the Larmor fluctuations at large N ?	559
5.3.4	Optimization of squeezing over time	559
5.3.4.1	What can be expected	559
5.3.4.2	Analysis at the first time scale	560
5.3.4.3	Analysis at the second time scale	561
5.3.4.4	From one regime to another: competition between noise and the thermodynamic limit	563
5.4	Implementation in Bose condensed atomic gases in the two-mode approximation	564
5.4.1	Nonlinear dynamics in the homogeneous case	566
5.4.2	Nonlinear dynamics in the trapped case	568

5.4.2.1	For stationary ϕ_σ modes	568
5.4.2.2	For unstationary ϕ_σ modes, difficulties and way out	571
5.4.3	Physical consequences of model Hamiltonians (5.114) and (5.127) and N fluctuations	572
5.5	Realistic and multimode study of the limits of spin squeezing in Bose condensed atomic gases	575
5.5.1	Simple considerations and effect of losses	575
5.5.1.1	Motivations and a heuristic calculation of the squeezing limit with the phase operators $\hat{\theta}_\sigma$	575
5.5.1.2	Effect of particle losses in the case $g_{aa} = g_{bb}$ and $g_{ab} = 0$	578
5.5.2	The proper Bogoliubov multimode study	584
5.5.2.1	Description of the experimental setup and procedure	584
5.5.2.2	How to carry out the calculation: contextualization of the Bogoliubov method, evolution equation of the variables $\hat{b}_{\sigma k}$, $\hat{b}_{\sigma k}^\dagger$ and $\hat{\theta}_{\sigma\phi}$ and their values at $t = 0^+$	587
5.5.2.3	Observables at $t > 0$ participating in spin squeezing	593
5.5.2.4	The final result on the spin squeezing factor	603
5.5.2.5	What happens at long times $t > 1/\gamma_{\text{coll}}$? Ergodic Limit	605
5.5.3	Complement: Effect of Branching the Interaction on a Condensed Gas	611
5.5.4	Application of Multimode Theory to the Spatially Homogeneous Case	613
5.5.4.1	Squeezing Optimum	613
5.5.4.2	Time Dependence of the Squeezing Factor	616
5.5.5	Application of Multimode Theory to the Trapped Case: The Limit of Spin Squeezing in an Isotropic Harmonic Trap with WKB	622
5.5.5.1	Motivation, Problem Formulation, Macroscopic Limit	622
5.5.5.2	WKB Approximation for Bogoliubov Modes	625
5.5.5.3	Classical Interpretation	627
5.5.5.4	The WKB Quantization Condition for energy levels	631
5.5.5.5	Justification of the reflection coefficient (–i) of WKB waves at turning points	633
5.5.5.6	Implementation of WKB in the calculation of ξ_{opt}^2	637
5.5.6	Complement: calculation of semiclassical integrals I, J, K .	641
5.5.6.1	Case $1 < 2j - 1 < \epsilon < j^2$: external trajectories	642
5.5.6.2	Case $0 < j^2 < \epsilon$: hybrid trajectories	643

5.5.7	Complement: calculation of χ_{Bog} , values of $\hat{D}$ and its variance, expression of $\hat{C}_{\text{osc}}(t)$ in the homogeneous case	644
5.5.7.1	Calculation of χ_{Bog}	645
5.5.7.2	A spin-off: value of the operator $\hat{D}$ and its variance	647
5.5.7.3	Another spin-off: expression of the operator $\hat{C}_{\text{osc}}(t)$ in the spatially homogeneous case	648
5.6	Schrödinger cat states and phase revival	650
5.6.1	Halfway between collapse and revival	650
5.6.2	Predictions of the Kitagawa-Ueda two-mode model	652
5.6.3	How to take advantage of the Schrödinger cat state in a clock experiment?	654
5.6.4	Effect of particle losses	656
5.6.4.1	Fidelity to the Schrödinger cat	657
5.6.4.2	Revival of the mean spin	659
5.6.5	Effect of a non-zero initial temperature: multimode analysis in the Bogoliubov approximation	662
5.6.5.1	Motivation	662
5.6.5.2	Proposed multimode protocol	662
5.6.5.3	Evolution over a single realization of the experiment	663
5.6.5.4	Fidelity to the spinorial Schrödinger cat	667
5.6.5.5	Constraint on temperature in an experiment	671
5.6.5.6	A nice underestimate	672
5.6.5.7	Fidelity to the orbito-spinorial Schrödinger cat	675
5.6.5.8	Revivals of the mean spin at $T \neq 0$	675
6	Temporal coherence of a condensate in an isolated gas: phase scrambling due to fluctuations in conserved quantities and phase diffusion due to interactions between quasiparticles	681
6.1	Introduction, motivations, overview, and measurability	681
6.1.1	A parallel between spatial and temporal coherence: condensation in spacetime	681
6.1.2	Objective and overview; scrambling versus diffusion	683
6.1.3	A Little-Studied Problem	686
6.1.4	A Measurable Coherence Function $g_1(t)$	687
6.2	Definition of the Problem, Its Reduction to Phase Dynamics, and Central Results on $g_1(t)$	688
6.2.1	The State of the System	688
6.2.2	Some simplifications: omission of fluctuations of $\hat{n}_\phi$, Gaussian approximation	689
6.2.3	Reduction to the microcanonical ensemble: correlation function of $d\hat{\theta}/dt$, phase shift variance, diffusion coefficient D and delay time t_0	691

6.2.4	Function $g_1(t)$ in the generalized statistical ensemble; Ballistic spreading with coefficient A	694
6.2.5	Explicit results on A , D , t_0 and the phase shift variance . .	696
6.2.5.1	Results on A	696
6.2.5.2	Results on D and t_0	698
6.2.6	Consequences on the spreading of the phase shift $\hat{\theta}(t) - \hat{\theta}(0)$	701
6.3	Calculating the Correlation Function $C(t)$ of $d\hat{\theta}/dt$	702
6.3.1	Overview	702
6.3.1.1	Everything rests on $C(t)$	702
6.3.1.2	Everything comes down to the microcanonical case	704
6.3.1.3	An obvious decorrelation... but false!	705
6.3.2	In the microcanonical ensemble by kinetic equations . . .	706
6.3.2.1	Statement of the problem	706
6.3.2.2	The kinetic equations	708
6.3.2.3	Their linearization, their stationary solution	709
6.3.2.4	The formal expression of $C_{mc}(\tau)$	711
6.3.2.5	In the subspace of zero angular momentum fluctuations	717
6.3.2.6	Time dependence of C_{mc}	718
6.3.2.7	Complement: behavior of $C_{mc}(\tau)$ when $\tau \rightarrow +\infty$.	720
6.3.3	In a generalized statistical ensemble with fixed N	723
6.3.3.1	What resembles the microcanonical case	723
6.3.3.2	What differs from the microcanonical case	724
6.3.4	The value of $C(+\infty)$: quantum ergodicity versus Markovian quantum master equation	727
6.3.4.1	By quantum ergodicity, i.e., microcanonicity of eigenstates	727
6.3.4.2	Failure of the Bogoliubov approximation on eigenstates	729
6.3.4.3	Failure of the Markovian quantum master equation	731
6.4	Further and Verification Studies of the Function $g_1(t)$ in the Microcanonical Ensemble	733
6.4.1	Numerical Classical Field Simulations	734
6.4.2	Resolvent Method	736
6.4.2.1	Position of the problem	736
6.4.2.2	Link between $g_{\theta,\lambda}(t)$ and a probability amplitude of presence in $ \psi_\lambda\rangle$	737
6.4.2.3	A simple reasoning using Fermi golden rule	738
6.4.2.4	Beyond Fermi golden rule: the resolvent method .	741
6.4.2.5	Phase diffusion results from the existence of a pole	745
6.4.3	Complement: Equality of the diffusion coefficients from the golden rule and the resolvent at extensively long times	745

7 A grand-canonical formulation of the Bogoliubov method and calculation of the ground state energy to Wu order	747
7.1 Introduction, motivation, and grand-canonical advantages	747
7.2 Hamiltonian model and expansion method	750
7.2.1 The grand canonical Hamiltonian of the lattice model . . .	750
7.2.2 Elimination of the condensate mode	751
7.2.2.1 A well-oiled procedure...	751
7.2.2.2 ... but what is the value of N ?	752
7.3 Order two in $f_{\text{nc}}^{1/2}$: Bogoliubov order	754
7.3.1 The quadratic Hamiltonian and its reduced form	754
7.3.2 The grand canonical equation of state and the noncondensed fraction	756
7.4 Order four in $f_{\text{nc}}^{1/2}$: Wu order for the ground energy level	760
7.4.1 Motivation	760
7.4.2 Interaction effective range correction at Bogoliubov order .	761
7.4.3 The grand canonical Hamiltonian at order $(f_{\text{nc}}^{1/2})^4$	763
7.4.4 Correction to the Bogoliubov grand potential at $T = 0$. . .	764
7.4.4.1 An application of perturbation theory	764
7.4.4.2 Taking the continuous limit $b/\xi \rightarrow 0$	768
7.4.4.3 Introduction of the three-body scattering hypervolume	769
7.4.4.4 Final result to order f_{nc}^2	772
7.4.4.5 Critical discussion of the result	773
7.4.5 Complement I: Correction $\Omega^{(4)}(\mu)$ to the Bogoliubov grand potential in the continuous limit $b/\xi \rightarrow 0$ of the lattice model	775
7.4.5.1 Convergent part C_{conv}^{Ω} , divergent part $J_{\mathcal{V}}(\epsilon)$	775
7.4.5.2 Study of $J_{\mathcal{V}}^{(1)}(\epsilon)$ for $\epsilon \rightarrow 0$	777
7.4.5.3 Study of $J_{\mathcal{V}}^{(2)}(\epsilon)$ for $\epsilon \rightarrow 0$	779
7.4.5.4 Conclusion on $\Omega^{(4)}(\mu)$	782
7.4.6 Complement II: Three-body scattering hypervolume D of the lattice model in the Born regime	782
7.4.6.1 The Model Hamiltonian	783
7.4.6.2 Form of the Scattering State: Only Unknown Function $f(\mathbf{r})$	783
7.4.6.3 A Closed Equation on $\hat{f}(\mathbf{k})$	785
7.4.6.4 Born Expansion of the Regular Part $\phi(\mathbf{k})$ of $\hat{f}(\mathbf{k})$. .	786
7.4.6.5 The sought result	789
7.4.7 Appendix to Complement II of Section 7.4.6	789
7.4.7.1 Expansion of $A(\mathbf{k}_1)$	789
7.4.7.2 Expansion of $B(\mathbf{k}_1)$	790
7.4.7.3 Expansion of $J(\mathbf{k}_1)$	790
7.4.7.4 Expansion of $K(\mathbf{k}_1)$	792

8 Case of reduced dimensionality: study of quasi-condensates using the Bogoliubov method in phase-modulus representation	795
8.1 Brief presentation and overview	795
8.1.1 More than a pale reflection of condensates, quasi-condensates	795
8.1.2 What theoretical angle of attack?	796
8.1.3 Directly connected to the experiments	797
8.2 Position of the problem and regime considered	799
8.2.1 At grand-canonical thermal equilibrium	799
8.2.2 Quasi-condensates under six conditions	800
8.3 Uncovering reduced dimensionality with the ordinary Bogoliubov method	802
8.3.1 Non-condensed density	803
8.3.1.1 Discussion by dimensionality	804
8.3.1.2 About Table 8.1	805
8.3.1.3 Moral on the coherence length	806
8.3.2 The pair distribution function	806
8.3.2.1 Quantum Contribution, Thermal Contribution	807
8.3.2.2 Moral on Density Fluctuations	808
8.3.3 Conclusion of the uncovering	809
8.4 Construction of the Model Hamiltonian	809
8.4.1 Scattering Length and Low-Energy T Matrix in Any Dimension	809
8.4.1.1 Motivation and Operational Definition	809
8.4.1.2 Low-Energy Zero-Range Model	811
8.4.1.3 Some Peculiarities of Dimension One	814
8.4.2 Lattice Model: Spacing, Coupling Constant, and Hamiltonian	816
8.4.2.1 The Model	816
8.4.2.2 The bare coupling constant g_0 via the T matrix	817
8.4.2.3 Constant g_0 , scattering length and effective range in 3D	819
8.4.2.4 Constant g_0 , scattering length and effective range in 2D	820
8.4.2.5 Constant g_0 , scattering length and effective range in 1D	822
8.4.2.6 The Hamiltonian in second quantization	823
8.4.2.7 How to choose the lattice spacing	825
8.5 Implementation of the phase-modulus Bogoliubov method	826
8.5.1 Switching to phase-modulus representation and idea of the method	826

8.5.2	Expansion of the Hamiltonian $\hat{H}_{\text{GC}}$ order by order in $\delta\hat{\rho}/\rho \approx b \mathbf{grad}\hat{\theta} \approx \epsilon$	829
8.5.2.1	Order 0 in ϵ	830
8.5.2.2	Order 1 in ϵ	831
8.5.2.3	Order 2 in ϵ	831
8.5.2.4	Order 3 in ϵ	832
8.5.3	Iterative solution up to order ϵ^2 and diagonalization of $\hat{H}_2$.	833
8.5.3.1	Order ϵ^0 : macroscopic wave function and nonlinear Schrödinger equation	833
8.5.3.2	Order ϵ^1 : nothing	834
8.5.3.3	Order ϵ^2 : fields $\hat{B}(\mathbf{r})$ and $\hat{\Lambda}_{\text{QC}}(\mathbf{r})$, collective variables $\hat{P}$ and $\hat{Q}$	835
8.5.3.4	Order ϵ^2 : Bogoliubov transformation, modal expansion, excitation spectrum, interpretation of $\hat{P}$ and $\hat{Q}$	837
8.5.3.5	In conclusion	841
8.6	Applications of Quasi-Condensate Theory	841
8.6.1	The Grand Potential Ω	841
8.6.2	Mean Density and Equation of State of the Gas	844
8.6.2.1	On the Need to Include a Cubic Correction	844
8.6.2.2	Direct Calculation of $\langle\delta\hat{\rho}\rangle_3$	845
8.6.2.3	Another, more elegant method	847
8.6.2.4	A third, more direct way	848
8.6.2.5	Equation of state: the results, their behavior at high and low temperatures	848
8.6.2.6	Thomas-Fermi Approximation in 2D	851
8.6.3	Pair Distribution Function g_2 : Formal Expression to Order ϵ^2	851
8.6.3.1	In Terms of the Field $\hat{\rho}(\mathbf{r})$	852
8.6.3.2	Skillful Transformations and Result	852
8.6.3.3	What variance of N ?	853
8.6.3.4	Relationship with ordinary Bogoliubov theory	854
8.6.4	First-order coherence function g_1 : formal expression to order ϵ^2	854
8.6.4.1	In terms of the fields $\hat{\rho}(\mathbf{r})$ and $\hat{\theta}(\mathbf{r})$	855
8.6.4.2	In the thermal state of $\hat{H}_2$	855
8.6.4.3	Correction due to $\hat{H}_3$	856
8.6.4.4	Synthesis and provisional form	859
8.6.4.5	An awkward factor $1/\rho_0$; final form	861
8.6.4.6	Overview of results; behavior at large distance	862
8.6.4.7	Wick's formula	865

8.6.5	Under what conditions are density fluctuations and phase gradient small?	866
8.6.5.1	Density fluctuations	867
8.6.5.2	Phase gradient	867
8.7	Detailed analysis of the correlation functions g_1 and g_2	868
8.7.1	In the thermodynamic limit for a continuous space	868
8.7.2	Long-distance study at $T = 0$	870
8.7.2.1	Obtaining asymptotic expansions	871
8.7.2.2	A simple physical application: fluctuations of the number of particles in a ball	875
8.7.3	Long-distance study at $T > 0$	877
8.7.3.1	Case of $g_1^{\text{Bog}}(\mathbf{r})$	877
8.7.3.2	Case of $g_2^{\text{QC}}(\mathbf{r})$	882
8.7.4	Short-distance study	887
8.7.4.1	Case of $g_2^{\text{QC}}(\mathbf{r})$; contact coefficient $\mathcal{C}$ and Hellmann-Feynman theorem	887
8.7.4.2	Case of $g_1^{\text{Bog}}(\mathbf{r})$; momentum distribution; Link with $\mathcal{C}$ and with the mean energy	893
8.8	Normal density, BKT transition, local or global superfluidity	898
8.8.1	Normal density	898
8.8.1.1	Definition	898
8.8.1.2	Expression to Bogoliubov order	899
8.8.1.3	Limiting behaviors with Temperature	900
8.8.1.4	A local normal density	901
8.8.2	The BKT transition	902
8.8.2.1	Summary presentation: universal discontinuity of $\rho_s(T)$, quantum vortices and activation energy	902
8.8.2.2	Transition temperature at Bogoliubov order: a rough estimate?	904
8.8.3	Moving quasi-condensates and global superfluidity	906
8.8.3.1	From the stationary case to the moving case	906
8.8.3.2	Probability of spontaneous overall motion and ρ_n	909
8.8.3.3	New normal densities: global ρ_n^{glob} , effective $\rho_n^{\text{eff}}(\nu)$ and differential $\rho_n^{\text{diff}}(\nu)$	910
8.8.3.4	Useful expression of ρ_n^{glob} and physical discussion	912
8.8.3.5	Seeing the overall motions	915
8.9	2D ground state energy beyond Bogoliubov-Popov	916
8.9.1	A simple application of chapter 7	916
8.9.2	Final form of the result and comparison to numerics	918
8.9.3	Application to the Thomas-Fermi limit	920
8.9.4	Effect of the interaction range	920

8.10 Complement: appearance of superfluid density ρ_s in the asymptotic behavior of $g_1(\mathbf{r}, \mathbf{r}')$ at $T > 0$	921
8.10.1 Problem statement and motivation	921
8.10.2 How to proceed	923
8.10.2.1 Logarithm, generating function and cumulants . .	923
8.10.2.2 Simplifying remarks	924
8.10.2.3 Coefficient B in $\phi(\mathbf{q})$ and form of the result	926
8.10.3 A calculation neglecting density fluctuations in function g_1	927
8.10.3.1 A clear physical motivation, a simple function $\phi_\theta(\mathbf{q})$	927
8.10.3.2 Second-order calculation in $\hat{H}_3$	929
8.10.3.3 First-order calculation in $\hat{H}_4$	932
8.10.3.4 Synthesis and coefficient B_θ	938
8.10.4 Asymptotic calculation on the full form of g_1	938
8.10.4.1 Formal expression of $\ln(g_1/\rho)$ to order ϵ^4	938
8.10.4.2 Calculation of averages in $\ln(g_1/\rho)$	940
8.10.4.3 Conclusion	945
Main Notations	947
Index	953
Bibliography	965